

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
6 May 2004 (06.05.2004)


PCT

(10) International Publication Number
WO 2004/038041 A1

- (51) International Patent Classification⁷: C12Q 1/48, G01N 33/569
- (21) International Application Number: PCT/GB2003/004592
- (22) International Filing Date: 22 October 2003 (22.10.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 0224997.7 26 October 2002 (26.10.2002) GB
- (71) Applicant (for all designated States except MG, US): ASTRAZENECA AB [SE/SE]; Sodertalje, S-151 85 (SE).
- (71) Applicant (for MG only): ASTRAZENECA UK LIMITED [GB/GB]; 15 Stanhope Gate, London, Greater London W1K 1LN (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): JAHIC, Haris [US/US]; AstraZeneca, 35 Gatehouse Drive, Waltham, MA 02451 (US). KERN, Gunther [DE/US]; AstraZeneca, 35 Gatehouse Drive, Waltham, MA 02451 (US).
- (74) Agent: ASTRAZENECA; Global Intellectual Property, P.O. Box 272, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4GR (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]

(54) Title: SCREENING ASSAY TO IDENTIFY INHIBITORS OF THE MURD ENZYME USING AN ACTIVATOR-INDEPENDENT MURD ENZYME

The lack of salt activation for *E. faecalis* MurD

(57) Abstract: The use of an activator-independent MurD enzyme in a screening assay to identify inhibitors of the enzyme, which assay comprises contacting the enzyme with a test compound in the presence of enzyme substrates and appropriate buffers and detecting any modulation of enzyme activity by the test compound.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

SCREENING ASSAY TO IDENTIFY INHIBITORS OF THE MURD ENZYME USING AN ACTIVATOR-INDEPENDENT MURD ENZYME

The present invention relates to improved screening assays and in particular to the use of activator-independent forms of the murein biosynthesis enzyme MurD, such as from
5 *Enterococcus faecalis* (*E. faecalis*). Such screening assays are used to identify and characterize modulators of the MurD enzyme.

Interest has been shown in the murein (Mur) biosynthesis pathway in bacteria, this is a key component of bacterial cell wall synthesis. Enzymes in this pathway are potential targets for broad-spectrum and selective antibacterial agents.

10 The bacterial enzyme MurD (UDP-N-acetylmuramyl-L-alanine:D-glutamate ligase catalyses the attachment of D-glutamate to a cytoplasmic peptidoglycan precursor, UDP-N-acetylmuramyl-L-alanine. This reaction results in the formation of a peptide linkage between the amino function of D-glutamate and the carboxyl terminus of UDP-N-acetylmuramyl-L-alanine. A stoichiometric consumption of ATP supplies the energy needed for this peptide
15 bond formation resulting in generation of ADP and orthophosphate.

Walsh et al (Journal of Bacteriology, Sept 1999, 181, No.17, 5395-5401) have examined the biochemical properties of the Mur D enzyme from two gram-negative bacteria, i.e. *Escherichia coli*, and *Haemophilus influenzae*, and two gram-positive bacteria i.e. *Enterococcus faecalis* and *Staphylococcus aureus*. They established data regarding the
20 biochemical properties of these enzymes and discussed similarities and differences between them, in particular with regard to salt-activation of the gram-negative bacteria. They report that the differences observed between the gram-positive and the gram-negative bacteria indicate that the two gram-negative bacteria may apply a more stringent regulation of cell wall biosynthesis at the early stage of the peptidoglycan biosynthesis pathway than do the two
25 gram-positive bacteria. However, the skilled reader is aware that the substrate purification procedure used by Walsh et al cannot remove all salts that may function as enzyme activators. Therefore it is not possible to draw meaningful conclusions as to the salt-dependence or lack of salt dependence of the two gram-positive bacteria.

Indeed, we have now found that the MurD enzyme from the gram-positive bacterium
30 *Staphylococcus aureus* is also salt-activated. This led to our discovery that the *E. faecalis* MurD enzyme has unique properties which make it possible to devise improved screening assays using an activator-independent MurD enzyme.

Therefore in a first aspect of the present invention, we provide the use of an activator-independent MurD enzyme in a screening assay to identify inhibitors of the enzyme, which assay comprises contacting the enzyme with a test compound in the presence of an enzyme substrate and appropriate buffers and detecting any modulation of enzyme activity by the test compound.

By "activator-independent" we mean that the Mur D enzyme is not activated by salt species normally associated with the substrate (here D-glutamic acid or more preferably UDP-N-acetylmuramyl-L-alanine) or other assay components. Monovalent cations such as ammonium (NH_4^+) and potassium (K^+) are particular salt species that activate Mur D. We note that if different amounts of substrate are used in an assay e.g. for K_m determinations or mode of inhibition studies, the amount of the activating cation such as ammonium ions is not constant. It can therefore not be clearly identified if an activity increase is due to activation or due to an increase of substrate concentration. Contrary to the results reported by Walsh et al. we have found that the *Staphylococcus aureus* Mur D enzyme is activated by cations such as NH_4^+ and K^+ (cf. Figure 2).

Our analysis shows that activator-dependent forms of the MurD enzyme have the following common amino acid residues i.e. G96, A112, A116, V126, L129, M133, G296, P298 and V422. The indicated positions are based on the *E. faecalis* Mur D sequence as set out in Figure 8 (and corresponding sequence alignments).

Therefore in a further aspect of the invention we provide the use of an activator-independent MurD enzyme which contains a MurD amino acid sequence wherein one or more of the amino acid residues at the positions given above is not as indicated for that particular amino acid. More conveniently, at least two, or at least three, or at least four, or at least five, or at least six, or at least seven, or at least 8, or all nine of the amino acid residues are not as indicated.

The activator-independent MurD may conveniently contain an amino acid sequence comprising one or more of the following residues i.e. K96, C112, G116, T126, M129, L133, N296, S298 and I422. More conveniently, at least two, or at least three, or at least four, or at least five, or at least six, or at least seven, or at least 8, or all nine of the amino acid residues are as indicated.

The activator-independent MurD enzyme conveniently comprises the following *E. faecalis* amino acid sequence

MKKITTYQNK KVLVLGLAKS GVSAAKLLHE LGALVTVNDA KQFDQNPDAQ
 DLLTLGIRVV TGGHPIELLD EEFELIVKNP GIPYTNPLVA EALTRKIPII
 TEVELAGQIA ECPIVGITGT NGKTTTTTMI GLLLNADRTA GEARLAGNIG
 FPASTVAQEA TAKDDLVMEL SSFQLMGIET FHPQIAVTN IFEAHLDYHG
 5 SRKEYVAAKW AIQKNMTVED TLILNWNQVE LQTLAKTTAA NVLPFSTKEA
 VEGAYLLDGK LYFNEEYIMP ADELGIPGSH NIENALAAIC VAKLKNVSNV
 QIRQTLKNFS GVPHRTQFVG EVQQRIFYND SKATNILATE MALSGFDNQK
 LLLLAGGLDR GNSFDELVPA LLGLKAIVLF GETKEKLAEA AKKANITIL
 FAENVQTAVT IAFDYSEKDD TILLSPACAS WDQYPNFEVR GEAFMQAVQQ
 10 LKESEM

or a Mur D amino acid sequence having at least 85% homology, such as 90 or 95% homology, therewith.

Alternatively the activator-independent Mur D amino acid sequence has at least 70% sequence identity with the above amino acid sequence.

15 Whilst we do not wish to be limited by theoretical considerations we believe that the activator-independent Mur D enzyme may have up to 16, such as up to 16, 14, 12, 10, 8, 6, 4, or 2, amino acids removed from the N-terminus and/or up to 12, such as up to 12, 10, 8, 6, 4, or 2, amino acids removed from the C-terminus of the enzyme.

Activator-independent Mur D sequences that do not correspond to published Mur D
 20 enzyme sequences are novel and represent a further aspect of the present invention.

Any convenient screening assay format may be used. By way of non-limiting example we disclose the following:

The enzyme substrates are conveniently UDP-MurNac-L-Ala, D-Glutamate and ATP.

25 The enzyme is conveniently pre-incubated with the test compound to allow inhibitors to bind to the enzyme. This may allow the detection of inhibitors with a slow binding mode to the enzyme or allow detection of specifically modifying inhibitors that may be out-competed by the substrates.

Any appropriate buffer can be used that has a pKa in the pH range where *E.faecalis* MurD is active (pH 7.0 – 10.0) Examples of convenient buffers include buffers that do not
 30 contain phosphate such as GOOD Buffer i.e. Tris or Hepes (Good, et al. (1966) Biochemistry, 5, 467-477). Modulation of enzyme activity may be detected using any convenient detection system, such as those which include a colour change eg. using malachite green. These include absorbance spectrophotometers, absorbance plate reader or any other instrument that

can determine the absorption of a solution between for example a wavelength range of 400 to 800 nm

Modulation of enzyme activity can be inhibition or activation of enzyme activity, conveniently enzyme inhibition.

5 Appropriate control reactions are conveniently performed to determine if a chemical compound interferes with the detection system and/or has an absorbance at the detection wavelength.

10 The test compound is any convenient compound that may be useful in pharmaceutical research. It may be a polypeptide of equal to or greater than, 2 amino acids such as up to 6 amino acids, up to 10 or 12 amino acids, up to 20 amino acids or greater than 20 amino acids such as up to 50 amino acids. For drug screening purposes, preferred compounds are chemical compounds of low molecular weight and potential therapeutic agents. They are, for example of molecular mass less than about 1000 Daltons, such as less than 800, 600 or 400 Daltons. If desired, the test compound may be a member of a chemical library. This may
15 comprise any convenient number of individual members, for example, tens to hundreds to thousands to millions etc., of suitable compounds, for example, peptides, peptoids and other oligomeric compounds (cyclic or linear), and template-based smaller molecules, for example, benzodiazepines, hydantoins, biaryls, carbocyclic and polycyclic compounds (eg. naphthalenes, phenothiazines, acridines, steroids etc.), carbohydrate and amino acids
20 derivatives, dihydropyridines, benzhydryls and heterocycles (eg. triazines, indoles, thiazolidines etc.). The numbers quoted and the types of compounds listed are illustrative, but not limiting. Preferred chemical libraries comprise chemical compounds of low molecular weight and potential therapeutic agents.

25 In a further aspect of the invention we provide a MurD enzyme modulator resulting from use of the assay method of the invention.

30 The activator-independent MurD enzyme may be produced using known recombinant techniques for cloning and expression (cf. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.). Convenient expression systems for the MurD enzyme include T7 promoter-driven transcription of the *murD* gene in a suitable host, more conveniently *E. coli*. Examples of convenient expression vectors include those with a T7 promoter and suitable cloning sites such as pET28b and pET30a (Novagen Inc. Madison WI USA). The *E. coli* host strains used for expression in such a system include those that contain the T7 RNA polymerase gene that

can be induced to initiate transcription of the *murD* gene, more specifically BL21(DE3) and HMS174(DE3).

The invention will now be illustrated by reference to the following Specific Description and Figures wherein:

5

Figure 1 shows the lack of salt activation for *E. faecalis* MurD and shows that all salts but SO₄-ions (=inhibitor) have no effect on *E. faecalis* MurD

Figure 2 shows the salt activation of *S. aureus* MurD

10

Figure 3 shows that the *E. coli* enzyme activity is dependent on the presence of ammonium formate while the *E. faecalis* enzyme activity is independent on the presence of this salt.

Figure 4 shows the stability of *E. faecalis* Mur D in DMSO at concentrations of up to 5%.

15

Figure 5 shows a graph (Eyring Plot) with the activation enthalpies for several MurD orthologues, including *E. faecalis*. The higher this value the more temperature dependent the reaction rate

20 Figure 6 (a) and (b) show the pH dependence of several MurD orthologues, including *E. faecalis*

Figure 7 shows the background ATPase activity of several MurD orthologues, including *E. faecalis*.

25

Specific Description:

We have established that the *E. faecalis* MurD enzyme has the following additional advantages for screening purposes compared to MurD enzymes from other species.

(a) it is not affected up to a dimethylsulfoxide (DMSO) concentration of 5%. Since DMSO is commonly used in screening and IC₅₀ measurements, this contributes to the stability of the assay results.

(b) its dependence of the reaction rate on temperature is low between 4°C to 40°C.

Temperature variations during assays have only a minor effect on the enzyme activity.

(c) it has the broadest pH optimum (pH 7.5 – 9.5). Changes in pH due to compound addition is less likely to affect activity and therefore assay results

(d) it has a low background ATPase activity. An idle ATPase activity can create a background signal in a screen (phosphate is being produced without catalysis occurring) so a low amount of this activity is desirable to produce a sensitive assay.

We devised a screening assay for this ortholog using Malachite Green for detection of the phosphate product. The assay includes controls to check for compound interference with the signal and interference with the detection method.

1. Compounds are preincubated with *E. faecalis* MurD for 5-30 min.

2. The 3 substrates ATP, UDP-MurNac-L-Ala and D-Glutamate are added to initiate the reaction.

3. After 30-60 min the reaction is stopped with a Malachite Green solution. The signal is recorded spectrophotometrically 4-10 min after the Malachite Green solution had been added. To control for compound interference with the assay signal, compounds are submitted to the same procedure but in the absence of *E. faecalis* MurD in step 1. Compounds interfering with the signal show an increased signal relative to a control where no *E. faecalis* MurD was present

To control for compound interference with the Malchite Green detection method, a constant amount of phosphate (10-15 uM) is replaced for *E. faecalis* MurD in step 1. Interference is detected by a increase or decrease of the signal relative to a control where no compound was present.

Specific assay conditions are 0.3nM *E. faecalis* MurD in 50mM Tris, 2.5mM DTT, 10mM MgCl₂, 0.01% Triton X-100, 50 uM ATP, 50uM UMA, 100uM D-Glu, pH 8.0. Typically

MurD is preincubated with inhibitor for 15 min in the absence of substrates. Subsequently the reaction is initiated by adding substrates and stopped by addition of Malachite Green after 60min. Signal is read 5 min after stopping the reaction. Although MurD activity is detected in this assay by formation of phosphate it is not limited to this detection method and can also be followed alternatively, for example, by measuring formation of other reaction products (UDP-N -acetylmuramyl-L-alanine-D-Glutamate, ADP) as well as the disappearance of the substrates (D-Glutamate, UDP-N-acetylmuramyl-L-alanine, ATP).

Claims:

1. The use of an activator-independent MurD enzyme in a screening assay to identify inhibitors of the enzyme, which assay comprises contacting the enzyme with a test compound in the presence of enzyme substrates and appropriate buffers and detecting any modulation of enzyme activity by the test compound.
2. The use as claimed in claim 1 wherein the activator-independent MurD enzyme comprises an amino acid sequence wherein one or more of the amino acid residues G96, A112, A116, V126, L129, M133, G296, P298 and V422 is a variant amino acid residue.
3. The use as claimed in claim 2 and wherein the activator-independent MurD enzyme comprises an amino acid sequence having one or more of the amino acid residues K96, C112, G116, T126, M129, L133, N296, S298 and I422.
4. The use as claimed in claim 1 and wherein the activator-independent MurD enzyme is an *E. faecalis* MurD enzyme.
5. The use as claimed in claim 1 and wherein the activator-independent MurD enzyme comprises the amino acid sequence
- MKKITTYQNK KVLVLGLAKS GVSAAKLLHE LGALVTVNDA KQFDQNPDAQ
DLLTLGIRVV TGGHPIELLD EEFELIVKNP GIPYTNPLVA EALTRKIPH
TEVELAGQIA ECPIVGITGT NGKTTTTTMI GLLLNADRTA GEARLAGNIG
FPASTVAQEA TAKDDLVMEL SSFQLMGIET FHPQIAVITN IFEAHLDYHG
SRKEYVAAKW AIQKNMTVED TLILNWNQVE LQTLAKTTAA NVLPFSTKEA
VEGAYLLDGK LYFNEEYIMP ADELGIPGSH NIENALAAIC VAKLKNVSNV
QIRQTLKNFS GVPHRTQFVG EVQQRREFYND SKATNILATE MALSGFDNQK
LLLLAGGLDR GNSFDELVPA LLGLKAIVLF GETKEKLAEA AKKANIETIL
FAENVQTAVT IAFDYSEKDD TILLSPACAS WDQYPNFEVR GEAFMQAVQQ
LKESEM or a sequence having at least 85% sequence identity therewith.

6. The use as claimed in any preceeding claim and wherein the activator-independent MurD enzyme has up to 16 amino acids removed from the N-terminus, or up to 12 amino acids removed from the C-terminus, of the amino acid sequence.

5 7. The use as claimed in any preceeding claim and wherein the enzyme substrates are UDP-MurNac-L-Ala, D-Glutamate and ATP.

8. The use as claimed in any preceeding claim and wherein the enzyme is pre-incubated with the test compound before contacting it with the substrates.

10

15

20

25

30

Figure 1

The lack of salt activation for *E. faecalis* MurD





Figure 2**Salt activation of *S. aureus* MurD**


Figure 3***E. coli* enzyme activity is dependent on ammonium formate**


Figure 4**DMSO dependence of *E. faecalis* MurD**

Figure 5**Temperature dependence of MurD orthologs**

This Graph (Eyring Plot) shows the activation enthalpies for all for orthologs. The higher this value the more temperature dependent the reaction rate.





Figure 6**pH dependence of MurD orthologs**

(a)



(b)





Figure 7**Background ATPase activity of MurD orthologs**

- 1 -

SEQUENCE LISTING

<110> AstraZeneca AB

5 <120> Method

<130> NGP/100874 Sep03

<150> GB 0224997.7

10 <151> 2002-10-26

<160> 4

<170> PatentIn version 3.2

15 <210> 1

<211> 463

<212> PRT

<213> Escherichia Coli bacterium

20

<220>

<221> misc_feature

<222> (1)..(3)

25 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (46)..(48)

30 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (73)..(73)

35 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (142)..(143)

40 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (154)..(155)

45 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (235)..(235)

50 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (247)..(249)

55 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (348)..(349)

60 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (399)..(400)

65 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

- 2 -

<222> (408)..(409)

<223> Xaa can be any naturally occurring amino acid

<220>

5 <221> misc_feature

<222> (460)..(463)

<223> Xaa can be any naturally occurring amino acid

<400> 1

10

Xaa Xaa Xaa Met Ala Asp Tyr Gln Gly Lys Asn Val Val Ile Ile Gly
 1 5 10 15

15

Leu Gly Leu Thr Gly Leu Ser Cys Val Asp Phe Phe Leu Ala Arg Gly
 20 25 30

20

Val Thr Pro Arg Val Met Asp Thr Arg Met Thr Pro Pro Xaa Xaa Xaa
 35 40 45

25

Gly Leu Asp Lys Leu Pro Glu Ala Val Glu Arg His Thr Gly Ser Leu
 50 55 60

30

Asn Asp Glu Trp Leu Met Ala Ala Xaa Asp Leu Ile Val Ala Ser Pro
 65 70 75 80

35

Ile Glu Ile Val Gly Asp Ile Glu Leu Phe Cys Arg Glu Ala Gln Ala
 100 105 110

40

Pro Ile Val Ala Ile Thr Gly Ser Asn Gly Lys Ser Thr Val Thr Thr
 115 120 125

45

Leu Val Gly Glu Met Ala Lys Ala Ala Gly Val Asn Val Xaa Xaa Gly
 130 135 140

50

Val Gly Gly Asn Ile Gly Leu Pro Ala Xaa Xaa Leu Met Leu Leu Asp
 145 150 155 160

Asp Glu Cys Glu Leu Tyr Val Leu Glu Leu Ser Ser Phe Gln Leu Glu
 165 170 175

55

Thr Thr Ser Ser Leu Gln Ala Val Ala Ala Thr Ile Leu Asn Val Thr
 180 185 190

60

Glu Asp His Met Asp Arg Tyr Pro Phe Gly Leu Gln Gln Tyr Arg Ala
 195 200 205

65

Ala Lys Leu Arg Ile Tyr Glu Asn Ala Lys Val Cys Val Val Asn Ala
 210 215 220

Asp Asp Ala Leu Thr Met Pro Ile Arg Gly Xaa Ala Asp Glu Arg Cys
 225 230 235 240

- 3 -

Val Ser Phe Gly Val Asn Xaa Xaa Xaa Met Gly Asp Tyr His Leu Asn
 245 250 255
 5
 His Gln Gln Gly Glu Thr Trp Leu Arg Val Lys Gly Glu Lys Val Leu
 260 265 270
 10
 Asn Val Lys Glu Met Lys Leu Ser Gly Gln His Asn Tyr Thr Asn Ala
 275 280 285
 15
 Leu Ala Ala Leu Ala Leu Ala Asp Ala Ala Gly Leu Pro Arg Ala Ser
 290 295 300
 20
 Ser Leu Lys Ala Leu Thr Thr Phe Thr Gly Leu Pro His Arg Phe Glu
 305 310 315 320
 Val Val Leu Glu His Asn Gly Val Arg Trp Ile Asn Asp Ser Lys Ala
 325 330 335
 25
 Thr Asn Val Gly Ser Thr Glu Ala Ala Leu Asn Xaa Xaa Gly Leu His
 340 345 350
 30
 Val Asp Gly Thr Leu His Leu Leu Leu Gly Gly Asp Gly Lys Ser Ala
 355 360 365
 35
 Asp Phe Ser Pro Leu Ala Arg Tyr Leu Asn Gly Asp Asn Val Arg Leu
 370 375 380
 40
 Tyr Cys Phe Gly Arg Asp Gly Ala Gln Leu Ala Ala Leu Arg Xaa Xaa
 385 390 395 400
 Pro Glu Val Ala Glu Gln Thr Xaa Xaa Glu Thr Met Glu Gln Ala Met
 405 410 415
 45
 Arg Leu Leu Ala Pro Arg Val Gln Pro Gly Asp Met Val Leu Leu Ser
 420 425 430
 50
 Pro Ala Cys Ala Ser Leu Asp Gln Phe Lys Asn Phe Glu Gln Arg Gly
 435 440 445
 55
 Asn Glu Phe Ala Arg Leu Ala Lys Glu Leu Gly Xaa Xaa Xaa Xaa
 450 455 460
 60
 <210> 2
 <211> 463
 <212> PRT
 <213> Escherichia faecalis bacterium
 65
 <220>
 <221> misc_feature
 <222> (200)..(200)
 <223> Xaa can be any naturally occurring amino acid

- 4 -

```

<220>
<221> misc_feature
<222> (261)..(262)
5 <223> Xaa can be any naturally occurring amino acid

<220>
<221> misc_feature
<222> (348)..(349)
10 <223> Xaa can be any naturally occurring amino acid

<220>
<221> misc_feature
<222> (365)..(365)
15 <223> Xaa can be any naturally occurring amino acid

<220>
<221> misc_feature
<222> (380)..(380)
20 <223> Xaa can be any naturally occurring amino acid

<400> 2

Met Lys Lys Ile Thr Thr Tyr Gln Asn Lys Lys Val Leu Val Leu Gly
25 1 5 10 15

Leu Ala Lys Ser Gly Val Ser Ala Ala Lys Leu Leu His Glu Leu Gly
30 20 25 30

Ala Leu Val Thr Val Asn Asp Ala Lys Gln Phe Asp Gln Asn Pro Asp
35 35 40 45

Ala Gln Asp Leu Leu Thr Leu Gly Ile Arg Val Val Thr Gly Gly His
40 50 55 60

Pro Ile Glu Leu Leu Asp Glu Glu Phe Glu Leu Ile Val Lys Asn Pro
45 65 70 75 80

Gly Ile Leu Tyr Thr Asn Pro Leu Val Ala Glu Ala Leu Thr Arg Lys
50 85 90 95

Ile Pro Ile Ile Thr Glu Val Glu Leu Ala Gly Gln Ile Ala Glu Cys
55 100 105 110

Pro Ile Val Gly Ile Thr Gly Thr Asn Gly Lys Thr Thr Thr Thr Thr
60 115 120 125

Met Ile Gly Leu Leu Leu Asn Ala Asp Arg Thr Ala Gly Glu Ala Arg
65 130 135 140

Leu Ala Gly Asn Ile Gly Phe Pro Ala Ser Thr Val Ala Gln Glu Ala
70 145 150 155 160

Thr Ala Lys Asp Asp Leu Val Met Glu Leu Ser Ser Phe Gln Leu Met
75 165 170 175

Gly Ile Glu Thr Phe His Pro Gln Ile Ala Val Ile Thr Asn Ile Phe

```

- 5 -

180 185 190
 5 Glu Ala His Leu Asp Tyr His Xaa Gly Ser Arg Lys Glu Tyr Val Ala
 195 200 205
 10 Ala Lys Trp Ala Ile Gln Lys Asn Met Thr Ala Glu Asp Thr Leu Ile
 210 215 220
 15 Leu Asn Trp Asn Gln Val Glu Leu Gln Thr Leu Ala Lys Thr Thr Ala
 225 230 235 240
 20 Ala Asn Val Leu Pro Phe Ser Thr Lys Glu Ala Val Glu Gly Ala Tyr
 245 250 255
 25 Leu Leu Asp Gly Xaa Xaa Lys Leu Tyr Phe Asn Glu Glu Tyr Ile Met
 260 265 270
 30 Pro Ala Asp Glu Leu Gly Ile Pro Gly Ser His Asn Ile Glu Asn Ala
 275 280 285
 35 Leu Ala Ala Ile Cys Val Pro Lys Leu Lys Asn Val Ser Asn Ala Gln
 290 295 300
 40 Ile Lys Gln Ser Leu Thr Asn Phe Ser Gly Val Pro His Arg Thr Gln
 305 310 315 320
 45 Phe Val Gly Glu Val Gln Gln Arg Arg Phe Tyr Asn Asp Ser Lys Ala
 325 330 335
 50 Thr Asn Phe Leu Ala Thr Glu Met Ala Leu Ser Xaa Xaa Gly Phe Asp
 340 345 350
 55 Asn Gln Lys Leu Leu Leu Leu Ala Gly Gly Leu Asp Xaa Arg Gly Asn
 355 360 365
 60 Ser Phe Asp Glu Leu Val Pro Ala Leu Leu Gly Xaa Leu Lys Ala Ile
 370 375 380
 65 Val Leu Phe Gly Glu Thr Lys Lys Lys Leu Ala Glu Ala Ala Lys Lys
 385 390 395 400
 70 Pro Asn Ile Glu Thr Ile Leu Phe Ala Glu Asn Val Gln Thr Ala Val
 405 410 415
 75 Thr Ile Ala Phe Asp Tyr Ser Glu Lys Asp Asp Thr Ile Leu Leu Ser
 420 425 430
 80 Pro Ala Cys Ala Ser Trp Asp Gln Tyr Pro Asn Phe Glu Val Arg Gly
 435 440 445
 85 Glu Ala Phe Met Gln Ala Val Gln Gln Leu Lys Glu Ser Glu Met

- 6 -

450

455

460

5 <210> 3
 <211> 463
 <212> PRT
 <213> Pseudomonas aeruginosa bacterium

10 <220>
 <221> misc_feature
 <222> (1)..(2)
 <223> Xaa can be any naturally occurring amino acid

15 <220>
 <221> misc_feature
 <222> (73)..(73)
 <223> Xaa can be any naturally occurring amino acid

20 <220>
 <221> misc_feature
 <222> (142)..(143)
 <223> Xaa can be any naturally occurring amino acid

25 <220>
 <221> misc_feature
 <222> (154)..(155)
 <223> Xaa can be any naturally occurring amino acid

30 <220>
 <221> misc_feature
 <222> (200)..(200)
 <223> Xaa can be any naturally occurring amino acid

35 <220>
 <221> misc_feature
 <222> (235)..(235)
 <223> Xaa can be any naturally occurring amino acid

40 <220>
 <221> misc_feature
 <222> (399)..(400)
 <223> Xaa can be any naturally occurring amino acid

45 <220>
 <221> misc_feature
 <222> (460)..(463)
 <223> Xaa can be any naturally occurring amino acid

50 <400> 3
 Xaa Xaa Met Ser Leu Ile Ala Ser Asp His Phe Arg Ile Val Val Gly
 1 5 10 15

55
 Leu Gly Lys Ser Gly Met Ser Leu Val Arg Tyr Leu Ala Arg Arg Gly
 20 25 30

60 Leu Pro Phe Ala Val Val Asp Thr Arg Glu Asn Pro Pro Glu Leu Ala
 35 40 45

65 Thr Leu Arg Ala Gln Tyr Pro Gln Val Glu Val Arg Cys Gly Glu Leu
 50 55 60

Asp Ala Glu Phe Leu Cys Ser Ala Xaa Arg Glu Leu Tyr Val Ser Pro

- 7 -

65 70 75 80
 5 Gly Leu Ser Leu Arg Thr Pro Ala Leu Val Gln Ala Ala Ala Lys Gly
 85 90 95
 10 Val Arg Ile Ser Gly Asp Ile Asp Leu Phe Ala Arg Glu Ala Lys Ala
 100 105 110
 15 Pro Ile Val Ala Ile Thr Gly Ser Asn Ala Lys Ser Thr Val Thr Thr
 115 120 125
 20 Leu Val Gly Glu Met Ala Val Ala Ala Asp Lys Arg Val Xaa Xaa Ala
 130 135 140
 25 Val Gly Gly Asn Leu Gly Thr Pro Ala Xaa Xaa Leu Asp Leu Leu Ala
 145 150 155 160
 30 Asp Asp Ile Glu Leu Tyr Val Leu Glu Leu Ser Ser Phe Gln Leu Glu
 165 170 175
 35 Thr Cys Asp Arg Leu Asn Ala Glu Val Ala Thr Val Leu Asn Val Ser
 180 185 190
 40 Glu Asp His Met Asp Arg Tyr Xaa Asp Gly Met Ala Asp Tyr His Leu
 195 200 205
 45 Ala Lys His Arg Ile Phe Arg Gly Ala Arg Gln Val Val Val Asn Arg
 210 215 220
 50 Ala Asp Ala Leu Thr Arg Pro Leu Ile Ala Xaa Asp Thr Val Pro Cys
 225 230 235 240
 55 Trp Ser Phe Gly Leu Asn Lys Pro Asp Phe Lys Ala Phe Gly Leu Ile
 245 250 255
 60 Glu Glu Asp Gly Gln Lys Trp Leu Ala Phe Gln Phe Asp Lys Leu Leu
 260 265 270
 65 Pro Val Gly Glu Leu Lys Ile Arg Gly Ala His Asn Tyr Ser Asn Ala
 275 280 285
 70 Leu Ala Ala Leu Ala Leu Gly His Ala Val Gly Leu Pro Phe Asp Ala
 290 295 300
 75 Met Leu Gly Ala Leu Lys Ala Phe Ser Gly Leu Ala His Arg Cys Gln
 305 310 315 320
 80 Trp Val Arg Glu Arg Gln Gly Val Ser Tyr Tyr Asp Asp Ser Lys Ala
 325 330 335
 Thr Asn Val Gly Ala Ala Leu Ala Ala Ile Glu Gly Leu Gly Ala Asp

- 8 -

340 345 350
 5 Ile Asp Gly Lys Leu Val Leu Leu Ala Gly Gly Asp Gly Lys Gly Ala
 355 360 365
 Asp Phe His Asp Leu Arg Glu Pro Val Ala Arg Phe Cys Arg Ala Val
 10 370 375 380
 Val Leu Leu Gly Arg Asp Ala Gly Leu Ile Ala Gln Ala Leu Xaa Xaa
 385 390 395 400
 15 Gly Asn Ala Val Pro Leu Val Arg Val Ala Thr Leu Asp Glu Ala Val
 405 410 415
 20 Arg Gln Ala Ala Glu Leu Ala Arg Glu Gly Asp Ala Val Leu Leu Ser
 420 425 430
 25 Pro Ala Cys Ala Ser Leu Asp Met Phe Lys Asn Phe Glu Glu Arg Gly
 435 440 445
 Arg Leu Phe Ala Lys Ala Val Glu Glu Leu Ala Xaa Xaa Xaa Xaa
 30 450 455 460
 <210> 4
 <211> 463
 <212> PRT
 35 <213> Staphylococcus aureus bacterium
 <220>
 <221> misc_feature
 40 <222> (73)..(73)
 <223> Xaa can be any naturally occurring amino acid
 <220>
 <221> misc_feature
 45 <222> (142)..(143)
 <223> Xaa can be any naturally occurring amino acid
 <220>
 <221> misc_feature
 50 <222> (200)..(200)
 <223> Xaa can be any naturally occurring amino acid
 <220>
 <221> misc_feature
 55 <222> (235)..(235)
 <223> Xaa can be any naturally occurring amino acid
 <220>
 <221> misc_feature
 60 <222> (261)..(262)
 <223> Xaa can be any naturally occurring amino acid
 <220>
 <221> misc_feature
 65 <222> (348)..(349)
 <223> Xaa can be any naturally occurring amino acid
 <220>

- 9 -

<221> misc_feature
 <222> (353)..(353)
 <223> Xaa can be any naturally occurring amino acid

5 <220>
 <221> misc_feature
 <222> (365)..(365)
 <223> Xaa can be any naturally occurring amino acid

10 <220>
 <221> misc_feature
 <222> (380)..(380)
 <223> Xaa can be any naturally occurring amino acid

15 <220>
 <221> misc_feature
 <222> (401)..(401)
 <223> Xaa can be any naturally occurring amino acid

20 <220>
 <221> misc_feature
 <222> (463)..(463)
 <223> Xaa can be any naturally occurring amino acid

25 <400> 4

Met Leu Asn Tyr Thr Gly Leu Glu Asn Lys Asn Val Leu Val Val Gly
 1 5 10 15

30 Leu Ala Lys Ser Gly Tyr Glu Ala Ala Lys Leu Leu Ser Lys Leu Gly
 20 25 30

35 Ala Asn Val Thr Val Asn Asp Gly Lys Asp Leu Ser Gln Asp Ala His
 35 40 45

40 Ala Lys Asp Leu Glu Ser Met Gly Ile Ser Val Val Ser Gly Ser His
 50 55 60

45 Pro Leu Thr Leu Leu Asp Asn Asn Xaa Pro Ile Ile Val Lys Asn Pro
 65 70 75 80

Gly Ile Pro Tyr Thr Val Ser Ile Ile Asp Glu Ala Val Lys Arg Gly
 85 90 95

50 Leu Lys Ile Leu Thr Glu Val Glu Leu Ser Tyr Leu Ile Ser Glu Ala
 100 105 110

55 Pro Ile Ile Ala Val Thr Gly Thr Asn Gly Lys Thr Thr Val Thr Ser
 115 120 125

60 Leu Ile Gly Asp Met Phe Lys Lys Ser Arg Leu Thr Gly Xaa Xaa Arg
 130 135 140

65 Leu Ser Gly Asn Ile Gly Tyr Val Ala Ser Lys Val Ala Gln Glu Val
 145 150 155 160

Lys Pro Thr Asp Tyr Leu Val Thr Glu Leu Ser Ser Phe Gln Leu Leu
 165 170 175

- 10 -

5 Gly Ile Glu Lys Tyr Lys Pro His Ile Ala Ile Ile Thr Asn Ile Tyr
 180 185 190
 Ser Ala His Leu Asp Tyr His Xaa Glu Asn Leu Glu Asn Tyr Gln Asn
 195 200 205
 10 Ala Lys Lys Gln Ile Tyr Lys Asn Gln Thr Glu Glu Asp Tyr Leu Ile
 210 215 220
 15 Cys Asn Tyr His Gln Arg Gln Val Ile Glu Xaa Ser Glu Glu Leu Lys
 225 230 235 240
 20 Ala Lys Thr Leu Tyr Phe Ser Thr Gln Gln Glu Val Asp Gly Ile Tyr
 245 250 255
 25 Ile Lys Asp Gly Xaa Xaa Phe Ile Val Tyr Lys Gly Val Arg Ile Ile
 260 265 270
 Asn Thr Glu Asp Leu Val Leu Pro Gly Glu His Asn Leu Glu Asn Ile
 275 280 285
 30 Leu Ala Ala Val Leu Ala Cys Ile Leu Ala Gly Val Pro Ile Lys Ala
 290 295 300
 35 Ile Ile Asp Ser Leu Thr Thr Phe Ser Gly Ile Glu His Arg Leu Gln
 305 310 315 320
 40 Tyr Val Gly Thr Asn Arg Thr Asn Lys Tyr Tyr Asn Asp Ser Lys Ala
 325 330 335
 45 Thr Asn Thr Leu Ala Thr Gln Phe Ala Leu Asn Xaa Xaa Ser Phe Asn
 340 345 350
 Xaa Gln Pro Ile Ile Trp Leu Cys Gly Gly Leu Asp Xaa Arg Gly Asn
 355 360 365
 50 Glu Phe Asp Glu Leu Ile Pro Tyr Met Glu Asn Xaa Val Arg Ala Met
 370 375 380
 55 Val Val Phe Gly Gln Thr Lys Ala Lys Phe Ala Lys Leu Gly Asn Ser
 385 390 395 400
 60 Xaa Gln Gly Lys Ser Val Ile Glu Ala Asn Asn Val Glu Asp Ala Val
 405 410 415
 65 Asp Lys Val Gln Asp Ile Ile Glu Pro Asn Asp Val Val Leu Leu Ser
 420 425 430
 Pro Ala Cys Ala Ser Trp Asp Gln Tyr Ser Thr Phe Glu Glu Arg Gly
 435 440 445

INTERNATIONAL SEARCH REPORT

PCT/GB 03/04592

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 C12Q1/48 G01N33/569

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WALSH A W ET AL: "Comparison of the D-glutamate-adding enzymes from selected gram-positive and gram-negative bacteria" JOURNAL OF BACTERIOLOGY, WASHINGTON, DC, US, vol. 181, no. 17, September 1999 (1999-09), pages 5395-5401, XP002249293 ISSN: 0021-9193 section Materials and Methods; p.5397, right-hand column, paragraph (i) and (ii).</p>	1-8
X	<p>WO 99 23241 A (EL SHERBEINI MOHAMMED ;MERCK & CO INC (US); GEISSLER WAYNE M (US);) 14 May 1999 (1999-05-14) abstract page 16, line 9 - line 29</p>	1,2

-/-

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

29 March 2004

Date of mailing of the international search report

06/04/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Jacques, P

INTERNATIONAL SEARCH REPORT

PCT/GB 03/04592

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 61050 A (MERCK & CO INC ;AZZOLINA BARBARA (US); EL SHERBEINI MOHAMED (US)) 2 December 1999 (1999-12-02) abstract page 17, line 14 - line 34 ---	1,2
A	PUCCI MJ ET AL: "Identification and characterization of cell wall-cell division gene clusters in pathogenic gram-positive cocci" JOURNAL OF BACTERIOLOGY, WASHINGTON, DC, US, vol. 179, no. 17, September 1997 (1997-09), pages 5632-5635, XP002098344 ISSN: 0021-9193 the whole document -----	1-8

INTERNATIONAL SEARCH REPORT

PCT/GB 03/04592

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9923241	A	14-05-1999	CA 2308620 A1	14-05-1999
			EP 1034297 A1	13-09-2000
			JP 2001521750 T	13-11-2001
			WO 9923241 A1	14-05-1999
			US 6534284 B1	18-03-2003
WO 9961050	A	02-12-1999	CA 2333667 A1	02-12-1999
			EP 1079855 A1	07-03-2001
			EP 1082875 A1	14-03-2001
			JP 2002516076 T	04-06-2002
			JP 2002516780 T	11-06-2002
			WO 9961050 A1	02-12-1999